arXiv:1210.7298v1 [math.GR] 27 Oct 2012

SEPARABILITY OF EMBEDDED SURFACES IN 3–MANIFOLDS

PIOTR PRZYTYCKI† AND DANIEL T. WISE‡

ABSTRACT. We prove that if S is a properly embedded π_1–injective surface in a compact 3–manifold M, then $\pi_1 S$ is separable in $\pi_1 M$.

1. INTRODUCTION

A subgroup $H \subset G$ is *separable* if H equals the intersection of finite index subgroups of G containing H. Scott proved that if $G = \pi_1 M$ for a manifold M with universal cover $\widetilde{M}$, then H is separable if and only if each compact subset of $H\backslash\widetilde{M}$ embeds in an intermediate finite cover of M [Sco78, Lem 1.4]. Thus, if $H = \pi_1 S$ for a compact surface $S \subset H\backslash\widetilde{M}$, then separability of H implies that S embeds in a finite cover of M. Rubinstein–Wang found a properly immersed π_1–injective surface $S \looparrowright M$ in a graph manifold such that S does not lift to an embedding in a finite cover of M, and they deduced that $\pi_1 S \subset \pi_1 M$ is not separable [RW98, Ex 2.6].

The objective of this paper is to prove:

Theorem 1.1. *Let M be a compact connected 3–manifold and let $S \subset M$ be a properly embedded connected π_1–injective surface. Then $\pi_1 S$ is separable in $\pi_1 M$.*

The problem of separability of an embedded surface subgroup was raised for instance by Silver–Williams — see [SW09] and the references therein to their earlier works. The Silver–Williams conjecture was resolved recently by Friedl–Vidussi in [FV12], who proved that $\pi_1 S$ can be separated from some element in $[\pi_1 M, \pi_1 M] - \pi_1 S$ whenever $\pi_1 S$ is not a fiber.

We proved Theorem 1.1 when M is a graph manifold in [PW11, Thm 1.1]. Theorem 1.1 was also proven when M is hyperbolic [Wis11]. In fact, every finitely generated subgroup of $\pi_1 M$ is separable for hyperbolic M, by [Wis11] in the case $\partial M \neq \emptyset$ and by Agol's theorem [Ago12] for M closed.

Overview: In Section 2 we introduce the basic notation and reduce to studying irreducible M that is *simple* in the sense that its Seifert-fibred components are

Date: October 30, 2012.

† Partially supported by MNiSW grant N201 012 32/0718 and the Foundation for Polish Science.

‡ Supported by NSERC.

products with base surfaces of sufficient complexity. In Section 3 we prove a topological result establishing separability of finite *semicovers* of M, i.e. maps required to be covers only over the interior of the blocks of the JSJ decomposition. This requires an omnipotence result for hyperbolic manifolds with boundary [Wis11, Cor 16.15] coming from virtual specialness.

To prove Theorem 1.1 we enhance the strategy employed in [PW11, Thm 1.1] for graph manifolds. Its main element was [PW11, Constr 4.13] which produced S*–injective* covers of M^g, which are covers $\overline{M^g}$ to which S lifts and, among other properties, such that the intersection with S is connected for each JSJ torus or component of $\overline{M^g}$. We extend the construction of S–injective semicovers to all compact 3–manifolds in Section 4. We use the double coset separability of relatively quasiconvex subgroups of π_1 of hyperbolic 3–manifolds with boundary [Wis11, Thm 16.23] and separability of double cosets of embedded surface subgroups of π_1 of graph manifolds [PW11, Thm 1.2].

We conclude with the proof of Theorem 1.1 in Section 5.

Acknowledgement: We thank Henry Wilton for discussions.

2. Framework and Reductions

Separability: We have the following finite index maneuverability: If $[H : H'] < \infty$ and $H' \subset G$ is separable, then $H \subset G$ is separable. Moreover, if $[G : G'] < \infty$ then a subgroup $H' \subset G'$ is separable if and only if $H' \subset G$ is separable. Finally, $H \subset G$ is separable if and only if for each $g \in G - H$ there is a finite quotient $\phi\colon G \to F$ with $\phi(g) \notin \phi(H)$. Thus G is *residually finite* when $\{1_G\}$ is separable. We will freely employ these statements.

Assumptions on M and S: Throughout this article M is a compact connected 3–manifold and might have nonempty boundary. We will make additional assumptions arising from the following reductions:

We can assume that S is not a sphere or a disc, since otherwise Theorem 1.1 follows from Hempel's residual finiteness of Haken 3–manifolds [Hem87] and Perelman's hyperbolization. By passing to a double cover we can assume that M is oriented. Furthermore, if S is not orientable, then the boundary $\widehat{S}$ of its tubular neighborhood is an oriented π_1–injective surface. As $[\pi_1 S : \pi_1 \widehat{S}] = 2$, the separability of $\pi_1 \widehat{S}$ implies separability of $\pi_1 S$. Hence we can assume that S is oriented. In the presence of our assumptions, the π_1–injectivity of S is equivalent to saying that S is *incompressible* and we will stay with this term.

Decomposition of M into blocks: An incompressible surface S in a reducible manifold can be homotoped into one of its prime factors, say M_0. Observe that there is a retraction $\pi_1 M \to \pi_1 M_0$ that kills the other factors. Consequently, if $g \in \pi_1 M_0 - \pi_1 S$, and we can separate g from $\pi_1 S$ in a finite quotient of $\pi_1 M_0$, then we can separate g from $\pi_1 S$ in a finite quotient of $\pi_1 M$. If $g \in \pi_1 M - \pi_1 M_0$,

then applying [Hem87] to the factors we can find a finite cover M' of M where all the terms of the normal form of g lie outside factor subgroups. Then the path representing g is nontrivial in the graph dual to the prime decomposition of M', and it suffices to use the residual finiteness of free groups. Hence we can assume that M is irreducible (though possibly ∂–reducible).

We will employ the *JSJ decomposition* of M, which is the minimal collection of incompressible tori (up to isotopy) each of whose complementary components is Seifert-fibred or atoroidal. If M is a single Seifert-fibred manifold, then all finitely generated subgroups of $\pi_1 M$ are separable [Sco78], so we can assume that M is not Seifert-fibred.

By passing to a double cover we can assume that there are no π_1–injective Klein bottles in M. We can also assume that M is not a torus bundle over the circle, since then the only embedded surfaces are the fibers. Now a complementary component of JSJ tori cannot be simultaneously Seifert-fibred and algebraically atoroidal. Algebraically atoroidal components are *hyperbolic* by hyperbolization, in other words, their interior carries a geometrically finite hyperbolic structure (possibly of infinite volume if there are non-toroidal boundary components, as in a handlebody). We will call these complementary components *hyperbolic blocks*. The other complementary components are Seifert-fibred and we assemble adjacent Seifert-fibred components into *graph manifold blocks*. The JSJ tori that are adjacent to at least one hyperbolic block are called *transitional*. An incompressible surface can be homotoped so that its intersection with each block is incompressible. Moreover, we can assume that S intersects each Seifert-fibred component along a horizontal or vertical surface, unless S is a ∂–parallel annulus. In the latter case separability follows easily from separability of the boundary torus (since it corresponds to a maximal abelian subgroup) and from a variant of Lemma 3.1 with T^* in the boundary.

The m–characteristic covers and simplicity: For a manifold E let $E_{[m]}$ denote the *m–characteristic* cover of E, which is the regular cover corresponding to the intersection of all subgroups of index m in $\pi_1 E$. In particular, if T is a torus, then $T_{[m]}$ is the cover corresponding to the subgroup $m\mathbb{Z} \times m\mathbb{Z} \subset \mathbb{Z} \times \mathbb{Z} = \pi_1 T$. A Seifert-fibred manifold E is *simple* if it is the product of the circle with a surface of genus ≥ 1 that has at least 2 boundary components. This boundary hypothesis (not required in [PW11]) ensures that there is a retraction onto each boundary component. Consequently, $E_{[m]}$ restricts to m–characteristic covers on boundary tori. An irreducible 3–manifold M is *simple* if its Seifert-fibred components are simple. We will pass to a simple finite cover of M in Lemma 3.1.

3. Extending semicovers to covers

We begin this section with the following additional simplification:

Lemma 3.1. *Let M be an irreducible 3–manifold that is not Seifert-fibred. Then M has a finite cover M' that is simple. Moreover, given covers $\{T^*\}$ of the transitional tori $\{T\}$ in M, we can assume that all the tori of M' covering T are isomorphic and factor through T^*.*

A key element of the proof employs the following omnipotence result for hyperbolic 3–manifolds with boundary.

Lemma 3.2 ([Wis11, Cor 16.15])**.** *Let M^h be a hyperbolic 3–manifold with boundary tori $\{T\}$. There exist finite covers $\{\widehat{T}\}$ such that for any further finite covers $\{T'\}$ there exists a finite cover $M^{h'}$ of M^h that restricts on boundary tori to $\{T'\}$.*

By passing to a further cover we can assume that $M^{h'} \to M^h$ is regular.

Proof of Lemma 3.1. Luecke and Wu proved in [LW97, Prop 4.4] that every graph manifold block M^g of M has a finite cover $M^{g'}$ that is simple. Without loss of generality we can assume that $M^{g'} \to M^g$ is regular.

Choose m such that

(i) for any M^g adjacent along a torus T to a hyperbolic block M^h, the cover $T'_{[m]}$ of the torus $T' \subset \partial M^{g'}$ covering T factors through $\widehat{T}$ of Lemma 3.2 and through T^*.

(ii) for a transitional or boundary torus $T \subset M$ adjacent to a hyperbolic block M^h but not to a graph manifold block, the cover $T_{[m]}$ factors through $\widehat{T}$ of Lemma 3.2 and through T^*, if T is transitional.

By Lemma 3.2, each hyperbolic block M^h of M has a finite regular cover $M^{h'}$ restricting on the boundary to $\{T'_{[m]}\}$ of (i) or $\{T_{[m]}\}$ of (ii). For a Seifert-fibred component E of one of the simple graph manifolds $M^{g'}$, as E is simple its retractive property ensures that the cover $E_{[m]}$ restricts to m–characteristic covers on its boundary tori. Gluing appropriately many copies of the various $E_{[m]}$ and $M^{h'}$ together provides the desired simple cover M' of M. □

Henceforth we **always** assume that M is simple (and irreducible as assumed in Section 2).

Definition 3.3. A *semicover* $\overline{M}$ of M is a local embedding $\overline{M} \to M$ that restricts to a covering map over each transitional torus and over each open block. Thus $\overline{M}$ can only fail to be a covering map at a component of $\partial\overline{M}$ that covers a transitional torus $T \subset M$. We say that $\overline{M} \to M$ is *finite* if $\overline{M}$ is compact.

We can now prove the main result of this section.

Proposition 3.4. *Any finite semicover $\overline{M}$ of M has a finite cover $\overline{M}' \to \overline{M}$ that embeds in a finite cover M' of M.*

Proof of Proposition 3.4. By Lemma 3.1, there is a cover $\widehat{M}$ of M such that for each transitional torus T of M all of the tori $\widehat{T} \subset \widehat{M}$ covering T are isomorphic and factor through all the covers of T in $\overline{M}$.

Let $\overline{M}' \to \overline{M}$ be the pullback of the cover $\widehat{M} \to M$ via the semicover $\overline{M} \to M$. Then on all of its boundary tori the semicover $\overline{M'} \to M$ restricts to the corresponding tori in $\widehat{M}$. Gluing $\overline{M'}$ with appropriately many copies of the components of $\widehat{M}$ extends $\overline{M'}$ to a cover M' of M. □

4. Surface-injective semicovers

In this section we construct a family of semicovers of M to which a given surface $S \subset M$ lifts. We keep the assumptions from Section 2.

We will use the following case of a theorem of Martínez-Pedroza:

Theorem 4.1 ([MP09, Thm 1.1])**.** *Let $S_0 \subset M^h$ be an incompressible geometrically finite surface properly embedded in a hyperbolic manifold M^h. Let $\partial S_0 = C_1 \sqcup \ldots \sqcup C_k$ and suppose these circles are contained in boundary tori $T_1, \ldots, T_k$ of M^h (some T_i may coincide). Then for almost all cyclic covers T_i' of T_i to which C_i lift, the graph of spaces obtained by amalgamating S_0 with T_i' along C_i maps π_1–injectively into M^h and the image of its π_1 in $\pi_1 M^h$ is relatively quasiconvex.*

The separability of double cosets of relatively quasiconvex subgroups of π_1 of a hyperbolic 3–manifold with boundary was established in [Wis11, Thm 16.23]. Consequently, we have:

Corollary 4.2. *For almost all cyclic covers T_i' described in Theorem 4.1, the group $\pi_1(S_0 \sqcup_{\{C_i\}} \{T_i'\})$ is separable in $\pi_1 M^h$.*

Corollary 4.3. *The subgroup $\pi_1 S_0$ as well as the double cosets $\pi_1 S_0 \pi_1 T_i$ are separable in $\pi_1 M^h$.*

Definition 4.4. Let $S \subset M$ be an incompressible surface. A semicover $\overline{M} \to M$ to which S lifts is *S–injective* if for each hyperbolic or graph manifold block $\overline{B}$ of $\overline{M}$ the intersection $S \cap \overline{B}$ is connected. We allow S itself to be disconnected.

Lemma 4.5 ([PW11, Constr 4.13])**.** *Let $S \subset M^g$ be a possibly disconnected incompressible surface in a graph manifold. Suppose n is an integer divisible by all of the degrees of (possibly disconnected) covers $S \cap E \to F$, where $E \subset M^g$ is a Seifert-fibred component with base surface F, and $S \cap E$ is horizontal. Then there is a finite cover $\overline{M^g}$ of M^g to which S lifts such that for each torus $\overline{T} \subset \partial \overline{M^g}$ intersecting S:*

- *$S \cap \overline{T}$ is connected,*
- *$\overline{T}$ maps to a torus $T \subset \partial M^g$ with degree $\frac{n}{|S \cap T|}$.*

Here $|S \cap T|$ denotes the number of components in the intersection of the surface S with the torus T.

Proposition 4.6. *Let $S \subset M$ be an incompressible surface. Let S_0 be a component of intersection of S with a hyperbolic or graph manifold block M_0 of M. Let T_i be the (possibly repeating) tori of ∂M_0 intersected by S_0. Let $g \in \pi_1 M_0 - \pi_1 S_0$ (resp. $g_i \in \pi_1 M_0 - \pi_1 S_0 \pi_1 T_i$ for each i). Then there is a finite S–injective semicover $\overline{M}$ with $g \notin \pi_1 \overline{M_0}$ (resp. $g_i \notin \pi_1 \overline{M_0} \pi_1 T_i$), where $\overline{M_0}$ is the block of $\overline{M}$ containing the lift of S_0.*

To make sense of the double cosets $\pi_1 S_0 \pi_1 T_i$ inside $\pi_1 M_0$, pick basepoints x_i of M_0 in C_i and interpret $\pi_1 S_0, \pi_1 T_i$ as subgroups of $\pi_1 M_0$ determined by loops based at x_i staying in S_0, T_i, respectively.

Proof. In the case where we assume $g \notin \pi_1 S_0$, we use that $\pi_1 S_0$ is separable in $\pi_1 M_0$. If M_0 is hyperbolic, this follows from Corollary 4.3. If M_0 is a graph manifold, we use separability of embedded surfaces in graph manifolds [PW11, Thm 1.1]. Hence there is a finite cover $\overline{M_0^*} \to M_0$ to which S_0 lifts with $g \notin \pi_1 \overline{M_0^*}$.

In the case where we assume $g_i \notin \pi_1 S_0 \pi_1 T_i$ for all i, we use that each double coset $\pi_1 S_0 \pi_1 T_i$ is separable in $\pi_1 M_0$. This follows from Corollary 4.3 and [PW11, Thm 1.2]. Hence there exists a cover $\overline{M_0^*} \to M_0$ to which S_0 lifts with $g_i \notin \pi_1 \overline{M_0^*} \pi_1 T_i$. Let n_i be the degree of the restriction of $\overline{M_0^*} \to M_0$ to the torus intersecting (the lift of) S_0 along (the lift of) C_i.

Choose n so that it is divisible by the numbers in (a)–(c), and also satisfies (d):

(a) every $|S \cap T|$, where T is a transitional or boundary torus,
(b) the degrees of (possibly disconnected) covers $S \cap E \to F$, where $E \subset M$ is a Seifert-fibred component with base surface F, and $S \cap E$ is horizontal,
(c) each $n_i | S \cap T_i|$ as above.
(d) We also require $\frac{n}{|S \cap T|}$ to be the degree of one of the covers $T' \to T$ given by Theorem 4.1 for a geometrically finite component of $S \cap M^h$ in a hyperbolic block M^h of M.

We construct the semicover $\overline{M}$ in the following way. Start with a copy $\overline{S}$ of S. Let T be a transitional or boundary torus of M. For each component of $S \cap T$ we attach along the corresponding circle in $\overline{S}$ the degree $\frac{n}{|S \cap T|}$ cyclic cover $\overline{T}$ of T. The value $\frac{n}{|S \cap T|}$ is an integer by (a).

For each graph manifold block M^g of M consider the finite (possibly disconnected) cover $\overline{M^g}$ from Lemma 4.5 applied to the surface $S \cap M^g$. The boundary components of $\overline{M^g}$ intersecting S coincide with the $\overline{T}$ attached to $\overline{S}$ above.

Consider now a hyperbolic block M^h of M such that $S \cap M^h$ is a union of fibers. In this case we choose $\overline{M^h}$ to be the union of $|S \cap M^h|$ copies of degree $\frac{n}{|S \cap M^h|}$ cyclic covers of M^h to which components of $S \cap M^h$ lift. Again, components of $\partial \overline{M^h}$ coincide with $\overline{T}$, so that we can consistently attach the $\overline{M^h}$ to $\overline{S}$.

Finally, if $S \cap M^h$ is not a union of fibers, then π_1 of each of its components is relatively quasiconvex in $\pi_1 M^h$, so by (d) and Corollary 4.2, there is a finite cover $\overline{M^h}$ extending $S \cap M^h \cup \{\overline{T}\}$, and we consistently attach the $\overline{M}^h$ to $\overline{S}$.

At this point we have constructed a finite S–injective semicover $\overline{M}$, without yet separating g (resp. g_i). Now we replace the block $\overline{M_0}$ with its fiber product with $\overline{M_0^*}$. (Algebraically π_1 of the *fiber product* is $\pi_1\overline{M_0} \cap \pi_1\overline{M_0^*} \subset \pi_1 M_0$.) This is possible by (c) which guarantees that the fiber product agrees with $\overline{M_0}$ on its boundary components intersecting S_0. After this replacement, $\overline{M}$ satisfies the requirement on g (resp. g_i), by definition of $\overline{M_0^*}$. □

5. Separability

Proof of Theorem 1.1. Choose a basepoint of M in S outside all JSJ and boundary tori. Let $f \in \pi_1 M - \pi_1 S$. Consider the based cover M^S of M with fundamental group $\pi_1 S$. Let γ^S be a path in M^S starting at the basepoint and representing f. Then γ^S does not terminate on S. Assume that γ^S is chosen so that it does not *backtrack*, i.e. its image in M intersects the transitional tori a minimal number of times.

Firstly, consider the case where γ^S terminates in a block $M_0^S \subset M^S$ that intersects the lift of S. Denote $S_0 = S \cap M_0^S$ and let $M_0 \subset M$ be the block covered by M_0^S. In the case where S_0 contains the basepoint, let $g \in \pi_1 M_0$ be an element represented by a path in M_0^S from the basepoint to the endpoint of γ^S.

By Proposition 4.6 there is a finite S–injective semicover $\overline{M}$ of M with $g \notin \pi_1\overline{M_0}$. Thus γ^S projects to a path $\overline{\gamma}$ in $\overline{M}$ that ends in $\overline{M_0}$ outside the lift of S_0. By Proposition 3.4 the semicover $\overline{M}$ has a finite cover $\overline{M}'$ that extends to a finite cover M' of M. Since the endpoint of the lift of $\overline{\gamma}$ to M', which lies in $\overline{M}'$, does not terminate on the based connected component of the preimage of S, we have $f \notin \pi_1 M' \pi_1 S$, as desired.

Secondly, consider the case where γ^S terminates in a block of M^S disjoint from the lift of S. Let $T^S \subset M^S$ be then the first connected component of the preimage of a transitional torus $T \subset M$ crossed by γ^S and disjoint from S. Let M_0^S be the last block that γ^S travels through before it hits T^S. Let $S_0 = S \cap M_0^S$ and let $M_0 \subset M$ be the block covered by M_0^S. If T coincides with one of the tori $T_i \subset M_0$ crossed by S_0 along C_i, then let $x_i \in C_i$ be a basepoint for M_0. Let x_i' be a lift of x_i in T^S. We keep the notation x_i for the lift of x_i to $S_0 \subset M_0^S$. Let $g_i \in \pi_1 M_0$ be an element represented by a path in M_0^S from x_i to x_i'.

Since T^S is disjoint from S_0, we have $g_i \notin \pi_1 S_0 \pi_1 T_i$. By Proposition 4.6 there is a finite S–injective semicover $\overline{M}$ of M with $g_i \notin \pi_1\overline{M_0}\pi_1 T_i$ for all i. In other words, $\overline{\gamma}$ leaves $\overline{M_0}$ through a torus disjoint from S_0.

By Proposition 3.4 the semicover $\overline{M}$ has a finite cover $\overline{M}'$ that extends to a finite cover M' of M. By separability of the transitional tori groups (since they

are maximal abelian) and residual finiteness of the free group (dual to transitional tori), by replacing M' with a further cover we can assume that the lift of $\overline{\gamma}$ to M' does not pass twice through the same transitional torus.

Let $T' \subset M'$ be the projection of T^S. Consider the double cover M'' obtained by taking two copies of M', cutting along T', and regluing. Then the based connected component of the preimage of S lies in one copy of (the cut) M' in M'', while the endpoint of the lift of $\overline{\gamma}$ lies in the other copy. Hence $f \notin \pi_1 M'' \pi_1 S$, as desired. □

References

[Ago12] Ian Agol, *The virtual Haken conjecture*, with an Appendix by Ian Agol, Daniel Groves, and Jason Manning (2012), preprint, available at `arXiv:1204.2810`.

[HW08] Frédéric Haglund and Daniel T. Wise, *Special cube complexes*, Geom. Funct. Anal. **17** (2008), no. 5, 1551–1620.

[Hem87] John Hempel, *Residual finiteness for* 3*-manifolds*, Combinatorial group theory and topology (Alta, Utah, 1984), Ann. of Math. Stud., vol. 111, Princeton Univ. Press, Princeton, NJ, 1987, pp. 379–396.

[FV12] Stefan Friedl and Stefano Vidussi, *A vanishing theorem for twisted Alexander polynomials with applications to symplectic* 4*–manifolds* (2012), available at `arXiv:1205.2434`.

[LW97] John Luecke and Ying-Qing Wu, *Relative Euler number and finite covers of graph manifolds*, Geometric topology (Athens, GA, 1993), AMS/IP Stud. Adv. Math., vol. 2, Amer. Math. Soc., Providence, RI, 1997, pp. 80–103.

[MP09] Eduardo Martínez-Pedroza, *Combination of quasiconvex subgroups of relatively hyperbolic groups*, Groups Geom. Dyn. **3** (2009), no. 2, 317–342.

[PW11] Piotr Przytycki and Daniel T. Wise, *Graph manifolds with boundary are virtually special* (2011), submitted, available at `arXiv:1110.3513`.

[RW98] J. Hyam Rubinstein and Shicheng Wang, π_1*–injective surfaces in graph manifolds*, Comment. Math. Helv. **73** (1998), no. 4, 499–515.

[WS12] Daniel T. Wise and Michah Sageev, *Cores for quasiconvex actions* (2012), submitted, available at `http://www.math.mcgill.ca/wise/papers.html`.

[Sco78] Peter Scott, *Subgroups of surface groups are almost geometric*, J. London Math. Soc. (2) **17** (1978), no. 3, 555–565.

[SW09] Daniel S. Silver and Susan G. Williams, *Twisted Alexander polynomials and representation shifts*, Bull. Lond. Math. Soc. **41** (2009), no. 3, 535–540.

[Wis11] Daniel T. Wise, *The structure of groups with quasiconvex hierarchy* (2011), submitted, available at `http://www.math.mcgill.ca/wise/papers.html`.

Institute of Mathematics, Polish Academy of Sciences, Śniadeckich 8, 00-956 Warsaw, Poland,

E-mail address: `pprzytyc@mimuw.edu.pl`

Dept. of Math. & Stats., McGill Univ., Montreal, QC, Canada H3A 0B9

E-mail address: `wise@math.mcgill.ca`